

NON-EQUILIBRIUM REAL-TIME DYNAMICS OF QUANTUM FIELDS: LINEAR AND NON-LINEAR RELAXATION IN SCALAR AND GAUGE THEORIES

Daniel Boyanovsky^(a), Héctor J. de Vega^(b), Richard Holman^(c), S. Prem Kumar^(c), Robert D. Pisarski^(d) and Julien Salgado^(b)

(a) Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh PA. 15260, U.S.A

(b) LPTHE, Université Pierre et Marie Curie (Paris VI) et Denis Diderot (Paris VII), Tour 16, 1er. étage, 4, Place Jussieu, 75252 Paris, Cedex 05, France

(c) Department of Physics, Carnegie-Mellon University, Pittsburgh, PA. 15213, U.S.A.

(d) Department of Physics, Brookhaven National Laboratory, Upton, NY 11973, U.S.A.

Lecture delivered at TFT98, the Vth International Workshop on Thermal Field Theories and Their Applications, Ratisbona (Regensburg), Germany, 10-14 August 1998
(September 1998)

The real time evolution of field condensates is solved for small and large field amplitudes in scalar theories. For small amplitudes, the quantum equations of motion for the condensate can be **linearized** and solved by Laplace transform. The late time evolution turns to be determined by the singularities in the complex plane (one-particle poles, two- and multi-particle cuts, Landau cuts for non-zero initial temperature). In **hot** scalar electrodynamics, we solve the real time evolution of field condensates with soft length scales $\sim k^{-1} > (eT)^{-1}$. We rederive the HTL effective action using the techniques of non-equilibrium field theory for small amplitude condensates. We find that transverse gauge invariant condensates relax as $1/t^2$ and longitudinal condensates associated with plasmon (charge density) excitations relax with $1/[t \log^2 t]$ behavior to asymptotic amplitudes that are determined by the quasi-particle poles.

In the **nonlinear** regime (for large initial energy densities) we analyze the dynamics of dissipation and relaxation in the unbroken symmetry phase of scalar theory after **linear** instabilities (parametric) are shut-off by the quantum back-reaction. A new time scale emerges that separates the linear from the non-linear regimes. This scale is non-perturbative in the coupling and initial amplitude. The non-perturbative evolution is studied within the context of the $O(N)$ vector model in the large N limit. A combination of numerical analysis and the implementation of a dynamical renormalization group resummation via multi-time scale analysis reveals the presence of unstable bands in the nonlinear regime. These are associated with **power law** growth of quantum fluctuations, that result in power law relaxation and dissipation with **non-universal and non-perturbative dynamical anomalous exponents**.

I. INTRODUCTION

The dynamics of relaxation of inhomogeneous field configurations out of equilibrium is an important problem and a common theme in cosmology and high energy physics.

In high energy physics the experimental possibility of studying the chiral and quark-gluon phase transition with high luminosity hadron colliders and upcoming heavy-ion

colliders makes imperative the understanding of relaxation and transport processes in extreme environments.

In early cosmology, the energy and time scales during the inflationary stage of the universe calls for an out of equilibrium quantum field treatment. Moreover, the high energy densities involved ($\sim 1/g \sim 10^{12}$) make necessary the use of non-perturbative approaches as the large N and Hartree methods [1].

There is currently a great deal of interest in understanding non-perturbative real time dynamics in gauge theories at high temperature, both within the realm of heavy ion collisions and the study of the quark gluon plasma [4]- [6], as well as the possibility for anomalous baryon number violation in the electroweak theory. In both situations the dynamics of soft gauge fields with typical length scales $> (gT)^{-1}$ is non-perturbative.

Their treatment requires a resummation scheme where one can consistently integrate out the hard scales associated with momenta $\approx T$ to obtain an effective theory for the soft scales. This is the program of resummation of hard thermal loops [8]. Physically, the hard scale represents the typical energy of a particle in the plasma while the soft scale is associated with collective excitations [9].

Besides gauge theories, scalar models as the $O(4)$ sigma model are relevant to study the non-equilibrium relaxation of an inhomogeneous initial configuration due to quantum and thermal fluctuations. Collective states of pions as those in connection with DCC (Disordered Chiral Condensates) can be treated in such a way [13,15].

The main task to solve in this context is the Cauchy problem in quantum field theory. That is, given the initial quantum description at time $t = 0$, to find it for arbitrary time $t > 0$. By quantum description we mean the initial Fock state in case of pure states and the density matrix when we have a mixed quantum description.

The dynamics is determined as usual by the Fock space evolution equations

$$\begin{aligned} i \frac{\partial |\hat{\Psi}\rangle}{\partial t} &= H |\hat{\Psi}\rangle \quad \text{pure states} \\ i \frac{\partial \hat{\rho}}{\partial t} &= [\hat{\rho}, H] \quad \text{mixed states} . \end{aligned} \quad (1)$$

Here H is the second quantized field theoretic Hamiltonian of the system and $\hat{\rho}$ is the density matrix in the Fock space.

The expectation value of any operator $\mathcal{O}$ is given by

$$\langle \mathcal{O} \rangle = \text{Tr}[\hat{\rho}(t)\mathcal{O}] . \quad (2)$$

For thermal initial conditions with an initial temperature given by $1/\beta$, we choose the density matrix at $t = 0$ to be

$$\rho_i = e^{-\beta H_i} \quad (3)$$

where H_i is a free (quadratic) Hamiltonian. Notice that H_i is *not* the Hamiltonian of the system for $t > 0$. The system thus evolves out of equilibrium. The above expectation value can be rewritten easily as a functional integral defined on a complex-time contour. The contour has two branches running forward and backward in time and a third leg along the imaginary axis stretching to $t = -i\beta$. This is the standard Schwinger-Keldysh closed time path formulation of non-equilibrium field theory (see ref. [2,3,16] for details). Fields defined on the forward and backward time contours are accompanied with (+) and (−) superscripts respectively and are to be treated independently. The expectation value of any string of field operators may be obtained by introducing independent sources on the forward and backward time contours and taking functional derivatives of the generating functional with respect to these sources. The imaginary time leg of the complex time contour does not contribute to the dynamics.

II. THE AMPLITUDE EXPANSION IN SCALAR AND GAUGE THEORY

Let us consider a quantum scalar field $\phi(\mathbf{x}, t)$ coupled to itself and eventually to other fields. The order parameter is then defined by

$$\langle \phi(\mathbf{x}, t) \rangle = \text{Tr}[\hat{\rho}(t)\phi(\mathbf{x}, t)] . \quad (4)$$

For small values of the order parameter, we can linearize the evolution equations for the field $\phi(\mathbf{x}, t)$ with the result [10,16],

$$\ddot{\phi}(\mathbf{x}, t) - \nabla^2 \phi(\mathbf{x}, t) + M^2(T) \phi(\mathbf{x}, t) + \int d^3x' \int_0^t dt' \Sigma_{ret}(\mathbf{x} - \mathbf{x}', t - t') \phi(\mathbf{x}', t') \quad (5)$$

Here $\Sigma_{ret}(\mathbf{x} - \mathbf{x}', t - t')$ stands for the retarded self-energy of the field $\phi(\mathbf{x}, t)$ and for the renormalized mass including temperature corrections. Notice that eq.(5) respects *causality* since the field at time t only depends on the field values at earlier times $t' < t$.

To be more precise, small values of the order parameter for a self-coupled scalar field of mass M means that $\sqrt{\lambda}\phi \ll M$ where λ is the coupling constant.

In eq.(5), the self-energy kernel $\Sigma_{ret}(\mathbf{x} - \mathbf{x}', t - t')$ can be computed in perturbation theory to any finite order. Notice that the self-energy contains information about the initial state through the field propagators. Namely, perturbative calculations are carried out with the non-equilibrium scalar propagators

$$\langle \Phi^{(a)\dagger}(\mathbf{x}, t) \Phi^{(b)}(\mathbf{x}, t') \rangle = -i \int \frac{d^3k}{(2\pi)^3} G_k^{ab}(t, t') e^{-i\mathbf{k} \cdot (\mathbf{x} - \mathbf{x}')} ,$$

where $(a, b) \in \{+, -\}$.

$$\begin{aligned} G_k^{++}(t, t') &= G_k^>(t, t')\Theta(t - t') + G_k^<(t, t')\Theta(t' - t) , \\ G_k^{--}(t, t') &= G_k^>(t, t')\Theta(t' - t) + G_k^<(t, t')\Theta(t - t') , \\ G_k^{\pm\mp}(t, t') &= -G_k^{<(>)}(t, t') , \\ G_k^>(t, t') &= \frac{i}{2\omega_k} \left[(1 + n_k) e^{-i\omega_k(t-t')} + n_k e^{i\omega_k(t-t')} \right] , \\ G_k^<(t, t') &= \frac{i}{2\omega_k} \left[n_k e^{-i\omega_k(t-t')} + (1 + n_k) e^{i\omega_k(t-t')} \right] , \\ \omega_k &= \sqrt{\mathbf{k}^2 + m^2} \quad ; \quad n_k = \frac{1}{e^{\beta\omega_k} - 1} . \end{aligned}$$

Here $\beta \equiv 1/T_i$ and T_i is the initial temperature as defined above [eq.(3)].

The explicit expression and the properties of $\Sigma_{ret}(\mathbf{x} - \mathbf{x}', t - t')$ obviously depend on the fields coupled to $\phi(\mathbf{x}, t)$. Several cases with scalar and fermionic fields were discussed in ref. [10,16].

Let us consider now scalar quantum electrodynamics (SQED). In fact, our ultimate goal is to understand relaxational processes associated with off-shell effects, such as Landau damping, in a Non-Abelian gauge theory [5]. What we do is to treat the same problem in the context of scalar quantum electrodynamics (SQED) model [7]. To leading order, we expect that this should be a good analogue of what happens in the Non-Abelian case; it also has the advantage that it is simpler to deal with, and as we will see below, it can be cast from the outset in terms of *gauge invariant* variables. This will eliminate any ambiguities associated with the usual problem of gauge dependence of off-shell quantities.

We will start with some inhomogeneous field configuration which is excited in the SQED plasma at $t = 0$. What we want to do is to follow the time development of this configuration as it interacts with the hard modes in the plasma, and in particular, we want to know to determine how it relaxes in time.

We use a gauge invariant formulation of SQED. The SQED Lagrangian is given by

$$\begin{aligned} \mathcal{L} &= D_\mu \Phi^\dagger D^\mu \Phi - m^2 |\Phi|^2 - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} , \\ D_\mu \Phi &= (\partial_\mu - ieA_\mu) \Phi . \end{aligned} \quad (6)$$

We begin with the identification of the constraints associated with gauge invariance. The Abelian gauge theory has two first class constraints, namely Gauss's law

and vanishing canonical momentum for A_0 . What we will do is project the theory directly onto the physical Hilbert space, defined as usual as the set of states annihilated by the constraints. First, we obtain gauge invariant observables that commute with the first class constraints and write the Hamiltonian in terms of these. All of the matrix elements between gauge invariant states (annihilated by first class constraints) are the same as those that would be obtained by fixing Coulomb gauge $\nabla \cdot \mathbf{A} = 0$. The Hamiltonian in the physical subspace can be written solely in terms of transverse components and includes the instantaneous Coulomb interaction. The instantaneous Coulomb interaction can be traded for a *gauge invariant* Lagrange multiplier field $A_0(\mathbf{x}, t)$ (a non-propagating field whose canonical momentum is absent from the Hamiltonian) linearly coupled to the charge density $\rho(\mathbf{x}, t)$ and obeying the algebraic equation of motion $\nabla^2 A_0(\mathbf{x}, t) = \rho(\mathbf{x}, t)$. This leads to the following Lagrangian density:

$$\begin{aligned} \mathcal{L} = & \partial_\mu \Phi^\dagger \partial^\mu \Phi + \frac{1}{2} \partial_\mu \mathbf{A}_T \cdot \partial^\mu \mathbf{A}_T - e \mathbf{A}_T \cdot \mathbf{j}_T \\ & - e^2 \mathbf{A}_T \cdot \mathbf{A}_T \Phi^\dagger \Phi + \frac{1}{2} (\nabla A_0)^2 \\ & + e^2 A_0^2 \Phi^\dagger \Phi - i e A_0 (\Phi \dot{\Phi}^\dagger - \dot{\Phi}^\dagger \Phi) , \\ j_T = & i (\Phi^\dagger \nabla_T \Phi - \nabla_T \Phi^\dagger \Phi) . \end{aligned} \quad (7)$$

where A_T is the transverse component of the gauge field.

In order to provide an initial value problem for studying the relaxational dynamics of charge density fluctuations we introduce an external source $\mathcal{J}_L(\mathbf{x}, t)$ linearly coupled to A_0 and study the linear response to this perturbation. Furthermore, it is convenient to introduce external sources coupled to the transverse gauge fields to study the linear response of *transverse* gauge field configurations. These external fields could in principle play the role of a semiclassical configuration coupled to small perturbations in a linearized approximation. We thus take,

$$\mathcal{L} \rightarrow \mathcal{L} - \mathcal{J}_L(\mathbf{x}, t) A_0(\mathbf{x}, t) - \mathcal{J}_T(\mathbf{x}, t) \cdot \mathbf{A}_T(\mathbf{x}, t) . \quad (8)$$

The relaxational dynamics of our initial inhomogeneous configurations is clearly an out of equilibrium process, and needs to be treated by an appropriate formalism [2,3,16].

The non-equilibrium SQED Lagrangian is given by

$$\mathcal{L}_{noneq} = \mathcal{L} [\mathbf{A}_T^+, \Phi^+, \Phi^{\dagger+}, A_0^+] - \mathcal{L} [\mathbf{A}_T^-, \Phi^-, \Phi^{\dagger-}, A_0^-] .$$

Perturbative calculations are carried out with the scalar Green's functions given above in eq.(6) and the non-equilibrium photon propagators:

$$\langle A_{Ti}^{(a)}(\mathbf{x}, t) A_{Tj}^{(b)}(\mathbf{x}, t') \rangle = -i \int \frac{d^3 k}{(2\pi)^3} \mathcal{G}_{ij}^{ab}(k; t, t') e^{-i\mathbf{k} \cdot (\mathbf{x} - \mathbf{x}')} ,$$

$$\begin{aligned} \mathcal{G}_{ij}^{++}(k; t, t') &= \mathcal{P}_{ij}(\mathbf{k}) [\mathcal{G}_k^>(t, t') \Theta(t - t') + \mathcal{G}_k^<(t, t') \Theta(t' - t)] \\ \mathcal{G}_{ij}^{--}(k; t, t') &= \mathcal{P}_{ij}(\mathbf{k}) [\mathcal{G}_k^>(t, t') \Theta(t' - t) + \mathcal{G}_k^<(t, t') \Theta(t - t')] \\ \mathcal{G}_{ij}^{\pm\mp}(k; t, t') &= -\mathcal{P}_{ij}(\mathbf{k}) \mathcal{G}_k^{<(>)}(t, t') \\ \mathcal{G}_k^>(t, t') &= \frac{i}{2k} \left[(1 + N_k) e^{-ik(t-t')} + N_k e^{ik(t-t')} \right] , \\ \mathcal{G}_k^<(t, t') &= \frac{i}{2k} \left[N_k e^{-ik(t-t')} + (1 + N_k) e^{ik(t-t')} \right] , \\ N_k &= \frac{1}{e^{\beta k} - 1} . \end{aligned}$$

Here $\mathcal{P}_{ij}(\mathbf{k})$ is the transverse projection operator:

$$\mathcal{P}_{ij}(\mathbf{k}) = \delta_{ij} - \frac{k_i k_j}{k^2} . \quad (9)$$

With these tools we are ready to begin our analysis of non-equilibrium SQED.

The field evolution equations in SQED for the transverse gauge field take then the form

$$\begin{aligned} \ddot{A}_i^T(\mathbf{x}, t) - \nabla^2 A_i^T(\mathbf{x}, t) \\ + \int d^3 x' \int_0^t dt' \Sigma_{ret}(\mathbf{x} - \mathbf{x}', t - t')_{ij} A_j^T(\mathbf{x}', t') = 0 \end{aligned} \quad (10)$$

where $1 \leq i, j \leq 3$ and $\Sigma_{ret}(\mathbf{x} - \mathbf{x}', t - t')_{ij}$ stands for the transverse photon vacuum polarization. After calculation we find [7],

$$\begin{aligned} \Sigma_{\mathbf{k}}(s)_{ij} &= \delta_{ij} \frac{e^2 T^2}{6} - 2e^2 \int \frac{d^3 p}{(2\pi)^3 \omega_p \omega_{k+p}} \\ & \left[(1 + n_p + n_{p+k}) \frac{\omega_{k+p} + \omega_p}{s^2 + (\omega_{k+p} + \omega_p)^2} + \right. \\ & \left. + (n_p - n_{k+p}) \frac{\omega_{k+p} - \omega_p}{s^2 + (\omega_{k+p} - \omega_p)^2} \right] p_{Ti} p_{Tj} \end{aligned} \quad (11)$$

In the hard thermal loop limit (HTL) $T \gg k$, $T \gg m$ this yields,

$$\Sigma_{\mathbf{k}}(s)_{ij} = \frac{e^2 T^2}{12\pi} \int d\Omega \hat{p}_{Ti} \hat{p}_{Tj} \frac{s}{s + i\hat{\mathbf{p}} \cdot \mathbf{k}} \quad (12)$$

We see that the magnetic mass identically vanishes, as expected. Introducing the lightlike vector $\hat{\mathcal{E}} = (1, -\hat{\mathbf{p}})$ and recognizing that the s -variable may be replaced by ∂_t in the time domain, we obtained the HTL effective Lagrangian for the transverse electromagnetic field [7],

$$\delta \mathcal{L}_{trans} = -\frac{e^2 T^2}{6} \int \frac{d\Omega}{4\pi} A_{Ti}(\mathbf{x}, t) \hat{p}_{Ti} \left[\frac{\partial_0}{\hat{\mathcal{E}} \cdot \partial} \right] \hat{p}_{Tj} A_{Tj}(\mathbf{x}, t)$$

Therefore, the hard thermal loops are resummed by this method in a very straightforward way.

We got analogously for the longitudinal part of the gauge field [7],

$$\delta \mathcal{L}_{long} = -\frac{e^2 T^2}{6} \left[A_0^2(\mathbf{x}, t) + \int \frac{d\Omega}{4\pi} A_0(\mathbf{x}, t) \frac{\partial_0}{\hat{\mathcal{E}} \cdot \partial} A_0(\mathbf{x}, t) \right]$$

which is the known result and displays explicitly the Debye screening mass of charge density fluctuations.

III. EXACT FIELD TIME EVOLUTION BY LAPLACE TRANSFORM

The field evolution equations (5) and (10) in the amplitude approximation can be solved exactly by Fourier transform in space and Laplace transform in time [7,10,16]. For the scalar case we define,

$$\delta(\mathbf{k}, t) = \int \frac{d^3x}{(2\pi)^3} e^{-i\mathbf{k}\cdot\mathbf{x}} \phi(\mathbf{x}, t), \quad \tilde{\phi}_{\mathbf{k}}(s) = \int_0^\infty dt e^{-st} \delta(\mathbf{k}, t)$$

$$\Sigma_{\mathbf{k}}(s) = \int_0^\infty dt e^{-st} \int \frac{d^3x}{(2\pi)^3} e^{i\mathbf{p}\cdot\mathbf{x}} \Sigma(\mathbf{x}, t).$$

Notice that the Laplace transform variable s corresponds to i times the frequency ω . The evolution equations (5) can now be solved exactly,

$$\tilde{\phi}_{\mathbf{k}}(s) = \frac{s \delta(\mathbf{k}, 0) + \dot{\delta}(\mathbf{k}, 0)}{s^2 + k^2 + M^2(T) + \Sigma_{\mathbf{k}}(s)}, \quad (14)$$

We have now to perform the inverse Fourier and Laplace transforms:

$$\phi(\mathbf{x}, t) = \int d^3k e^{i\mathbf{k}\cdot\mathbf{x}} \int_{\Gamma} e^{st} \tilde{\phi}_{\mathbf{k}}(s) \frac{ds}{2\pi i} \quad (15)$$

where Γ is a contour running from $-i\infty$ to $+i\infty$ parallel to the imaginary s axis and with a small but nonzero real part. As it is obvious the singularities of eq.(14) are crucial for the behaviour of $\phi(\mathbf{x}, t)$. The zeroes of the denominator are specially important.

In order to evaluate (15) we deform the contour Γ to the left. Pole contributions are explicitly computed by the residue theorem. Cut contributions become real integrals of the corresponding cut discontinuities (see [7,10,16]).

In SQED we analogously obtain for the transverse gauge field,

$$A_{i\mathbf{k}}^T(t) = \int_{\Gamma} e^{st} \frac{ds}{2\pi i} \frac{s A_{i\mathbf{k}}^T(0) + \dot{A}_{i\mathbf{k}}^T(0) + \tilde{\mathcal{J}}_{Ti}(\mathbf{k}, s)}{s^2 + k^2 + \Sigma_{\mathbf{k}}^T(s)} \quad (16)$$

Here $\Sigma_{\mathbf{k}}^T(s)$ stands for the transverse vacuum polarization.

There are two kinds of cuts in $\Sigma_{\mathbf{k}}^T(s)$:

- the two particle production cuts from $s = \pm i(m + \omega_k)$ to $\pm i\infty$ where m is the mass of the scalar.
- the thermal (Landau) cut running from $-ik$ to $+ik$

It must be stressed that the Landau cut discontinuity is of order T^2 whereas the production cut discontinuity is of order $\ln T$. Hence, only the Landau cut will be considered in the HTL limit.

The explicit form for $\Sigma_{\mathbf{k}}^t(i\omega + 0^+)$ follows from eq.(11). In the HTL limit we find (see for example [7])

$$\Sigma_{\mathbf{k}}^t(i\omega + 0^+) = \frac{e^2 T^2}{12} \left[2 \frac{\omega^2}{k^2} + \frac{\omega}{k} \left(1 - \frac{\omega^2}{k^2} \right) \ln \left| \frac{k + \omega}{k - \omega} \right| \right]$$

$$+ i \frac{e^2 T^2 \pi \omega}{12 k} \left(1 - \frac{\omega^2}{k^2} \right) \Theta(k^2 - \omega^2) = \Sigma_{\mathbf{k}}^t(\omega)_R + i \Sigma_{\mathbf{k}}^t(\omega)_I$$

In addition to the branch cut singularities, the retarded propagator has isolated poles corresponding to the quasiparticle excitations which can propagate in the plasma. In the HTL limit they are at $s = \pm i\omega_P \simeq \pm i \frac{eT}{3}$.

After deforming the contour Γ we obtain [7]

$$\mathcal{A}_{Ti}(\mathbf{k}, t) = \mathcal{A}_{Ti}^{pole}(\mathbf{k}, t) + \mathcal{A}_{Ti}^{cut}(\mathbf{k}, t), \quad (17)$$

(13) The contributions from the quasiparticle poles add up to give a purely oscillatory behavior in time. The residues at the poles give rise to a T -dependent wave function renormalization. We thus obtain,

$$\mathcal{A}_{Ti}^{pole}(\mathbf{k}, t) = Z^t[T] \left[\mathcal{A}_{Ti}(\mathbf{k}, 0) \cos \omega_P t + \dot{\mathcal{A}}_{Ti}(\mathbf{k}, 0) \frac{\sin \omega_P t}{\omega_P} \right],$$

$$Z^t[T] = \left[1 - \frac{\partial \Sigma^t(i\omega)}{\partial \omega^2} \right]_{\omega=\omega_P \simeq eT/3}^{-1}. \quad (18)$$

Here $Z^t(T)$ is the temperature dependent wave function renormalization defined on-shell at the quasiparticle pole whose leading HTL contribution is obtained from the self-energy (17). The cut contribution is given by

$$\mathcal{A}_{Ti}^{cut}(\mathbf{k}, t) = \frac{2}{\pi} \int_0^k d\omega \Sigma_{Ti}^t(i\omega + 0^+)$$

$$\frac{\mathcal{A}_{Ti}(\mathbf{k}, 0) \omega \cos(\omega t) + \dot{\mathcal{A}}_{Ti}(\mathbf{k}, 0) \sin(\omega t)}{[\omega^2 - k^2 - \Sigma_{\mathbf{k}}^t(\omega)_R]^2 + [\Sigma_{\mathbf{k}}^t(\omega)_I]^2}. \quad (19)$$

Evaluating Eqs.(17),(18) and (19) at $t = 0$ we obtain an important sum rule,

$$Z^t[T] + \frac{2}{\pi} \int_0^k d\omega \frac{\omega \Sigma_{\mathbf{k}}^t(\omega)_I}{[\omega^2 - k^2 - \Sigma_{\mathbf{k}}^t(\omega)_R]^2 + [\Sigma_{\mathbf{k}}^t(\omega)_I]^2} = 1 \quad (20)$$

The integral over the cut (19) cannot be evaluated in closed form but its long time asymptotics is dominated by the end-point contribution from $\omega = k$. We find in the long time limit $t \gg 1/k$

$$\mathcal{A}_{Ti}^{cut}(\mathbf{k}, t) \stackrel{t \rightarrow \infty}{\simeq} -\frac{12}{e^2 T^2} \left\{ \mathcal{A}_{Ti}(\mathbf{k}, 0) \frac{\cos(kt)}{t^2} + \dot{\mathcal{A}}_{Ti}(\mathbf{k}, 0) \frac{\sin(kt)}{kt^2} \right\} \left[1 + \mathcal{O}\left(\frac{1}{t}\right) \right]. \quad (21)$$

We want to stress that the real time dynamics of the condensate is completely determined by the analytic structure of the retarded propagator and the **global structure** of the spectral density in the s -plane.

The second important point to note is that the long time behaviour is a **power law** $\sim t^{-2}$ (times oscillations) and **not an exponential decay**.

This means that Landau damping effects **cannot** be reproduced by phenomenological ‘viscous’ terms of the

type $\sim \Gamma \frac{d}{dt}$ neither at long nor at short times. The failure of such a phenomenologically motivated ansatz was already noticed at zero temperature in different contexts in ref. [3]. We stress that such a description not only fails to reproduce the power law behaviour but in fact ignores *all the non-local* physics of Landau damping which is so clearly encoded in the Hard Thermal Loop kernels.

A very similar analysis yields the longitudinal component of the gauge field in the case of an impulsive source $\mathcal{J}_L(\mathbf{x}, t) = \delta^3(\mathbf{x})\delta(t)$. The case of a more complicated source can be obtained by convolution. In this case we obtain [7]

$$\tilde{\mathcal{A}}_0(\mathbf{k}, s) = \frac{1}{k^2 + \Sigma^l(s)} \quad (22)$$

where in the HTL limit

$$\Sigma^l(s) = \frac{e^2 T^2}{3} - \frac{e^2 T^2}{6} \frac{s}{ik} \ln \left(\frac{is - k}{is + k} \right). \quad (23)$$

Thus the longitudinal self-energy along the imaginary axis in the s -plane, when approaching from the right is obtained as before to be:

$$\begin{aligned} \Sigma^l(i\omega + 0^+) &= \frac{e^2 T^2}{3} \left[1 - \frac{\omega}{2k} \ln \left| \frac{k + \omega}{k - \omega} \right| \right] \\ &\quad - i \frac{e^2 T^2 \pi}{6} \frac{\omega}{k} \Theta(k^2 - \omega^2). \end{aligned} \quad (24)$$

The location of the longitudinal quasiparticle or plasmon poles is given again for soft external momenta ($k \ll eT$) by $s = \pm i\omega_0 \simeq \pm ieT/3$. The real time dependence of the longitudinal condensate is then found by inverting the transform through

$$\mathcal{A}_0(\mathbf{k}, t) = \int_{c-i\infty}^{c+i\infty} \frac{ds}{2\pi i} e^{st} \tilde{\mathcal{A}}_0(\mathbf{k}, s) \quad (25)$$

where the contour is to the right of all the singularities as in Eq.(16). As in the transverse case, at high temperatures the singularity structure is dominated by the discontinuity across the cut that runs from $-ik$ to $+ik$ corresponding to Landau damping, and the two plasmon poles at $\pm i\omega_0$. Deforming the contour to pick up the cut and pole contributions we get

$$\mathcal{A}_0(\mathbf{k}, t) = \mathcal{A}_0^{pole}(\mathbf{k}, t) + \mathcal{A}_0^{cut}(\mathbf{k}, t), \quad (26)$$

where

$$\mathcal{A}_0^{pole}(\mathbf{k}, t) = -Z^l[T] \frac{\sin(\omega_0 t)}{\omega_0}, \quad Z^l[T] = \left[\frac{\partial \Sigma^l(i\omega)}{\partial \omega^2} \right]_{\omega=\omega_0 \simeq eT/3}^{-1} \quad (27)$$

and

$$\mathcal{A}_0^{cut}(\mathbf{k}, t) = -\frac{2}{\pi} \int_0^k d\omega \frac{\Sigma_k^l(\omega)_I \sin(\omega t)}{[k^2 + \Sigma_k^l(\omega)_R]^2 + [\Sigma_k^l(\omega)_I]^2}$$

Unlike the transverse components for which the sum rule is a consequence of the canonical commutation relations,

for the longitudinal component there is no equivalent sum rule because the field $A_0(x)$ is a non-propagating Lagrange multiplier.

The long time, asymptotic behaviour of the longitudinal condensate is dominated by the end-points of the integral (28). The end-point $\omega = 0$ yields contributions that vanish for long t faster than any negative power of t , as it was the case for the transverse part (19). We find that the long time asymptotics for $t \gg 1/k$ is dominated by the end-point contribution at $\omega = k$ [7],

$$\mathcal{A}_0^{cut}(\mathbf{k}, t) = a_{asy}^{cut}(k, t) \left[1 + \mathcal{O}\left(\frac{1}{t}\right) \right], \quad (28)$$

$$\begin{aligned} a_{asy}^{cut}(k, t) &\equiv -\frac{12}{e^2 T^2} \int_0^\infty e^{-x} dx \frac{\cos[kt + \alpha(x, t)]}{\sqrt{\log^4 \frac{cx}{kt} + \frac{5\pi^2}{2} \log^2 \frac{cx}{kt} + \frac{9\pi^4}{16}}} \\ \alpha(x, t) &\equiv \arctan \frac{\pi \log \frac{cx}{kt}}{\frac{3\pi^2}{4} + \log^2 \frac{cx}{kt}}, \quad c \equiv \frac{1}{2} \exp[2 + \frac{6k^2}{e^2 T^2}]. \end{aligned}$$

The integral in Eq.(28) cannot be expressed in terms of elementary functions and it is related to the $\nu(x)$ function [20]. For large t , one can derive an asymptotic expansion in inverse powers of $\log[kt/c]$ by integrating Eq.(28) by parts:

$$\begin{aligned} a_{asy}^{cut}(k, t) &\stackrel{t \rightarrow \infty}{\simeq} -\frac{12}{e^2 T^2} \frac{\cos(kt + \beta(kt))}{t \log^2 \frac{kt}{c} + \frac{\pi^2}{4}} \left[1 + \mathcal{O}\left(\frac{1}{\log t}\right) \right] \\ \beta(kt) &\equiv -\arctan \frac{\pi \log kt}{\log^2 kt - \frac{\pi^2}{4}}, \end{aligned} \quad (29)$$

This expansion is not very good quantitatively unless t is very large. For example, for $kt = 500$, $a_{asy}^{cut}(k, t)$ given by Eq.(28) approximates $\mathcal{A}_0^{cut}(\mathbf{k}, t)$ up to 0.1%, whereas, the dominant term in Eq.(29) is about 30% smaller than $a_{asy}^{cut}(k, t)$ [7].

To summarize, we gather the final results for the asymptotic real-time evolution of the transverse and longitudinal condensates in the linear approximation **(a) Transverse (no external source):**

$$\begin{aligned} \mathcal{A}_{Ti}(\mathbf{k}, t) &= \mathcal{A}_{Ti}(\mathbf{k}, 0) \left[Z^t[T] \cos \omega_p t - \frac{12}{e^2 T^2} \frac{\cos kt}{t^2} \right] \\ &\quad + \dot{\mathcal{A}}_{Ti}(\mathbf{k}, 0) \left[Z^t[T] \frac{\sin \omega_p t}{\omega_p} - \frac{12}{e^2 T^2} \frac{\sin kt}{kt^2} \right] + \mathcal{O}\left(\frac{1}{t^3}\right) \end{aligned}$$

The sum rule (20) implies that the coherent field configuration relaxes to an asymptotic amplitude which is *smaller* than the initial, and the ratio of the final to the initial amplitude is completely determined by the thermal wave function renormalization.

(b) Longitudinal (impulsive external source):

$$\mathcal{A}_0(\mathbf{k}, t) = -Z^l[T] \frac{\sin(\omega_0 t)}{\omega_0} + a_{asy}^{cut}(k, t) \left[1 + \mathcal{O}\left(\frac{1}{t}\right) \right].$$

The transverse and longitudinal wave function renormalizations $Z^t[T]$, $Z^l[T]$ are given by Eqs. (18) and (27) respectively and $a_{asy}^{cut}(k, t)$ is given by Eqs. (28)-(29) while

the positions of the poles ω_p , ω_0 can be obtained by the finding the zeroes of the denominators using eqs.(17) and (24), respectively.

In summary, we find that the long time dynamics is dominated by the Landau damping thresholds at $\omega = \pm k$, not by the $\omega \approx 0$ region of the spectral density. The early time dynamics is determined by moments of the total spectral density.

While an understanding of the real-time relaxation of non-equilibrium, inhomogeneous field configurations is of fundamental importance in the physics of relaxation in the QGP [5], such a calculation also has phenomenological implications for sphaleron induced B-violating processes.

One of our main focus is to assess in detail the real time non-equilibrium dynamics of soft excitations in the plasma with particular attention to a critical analysis of the long-standing belief that the small frequency region of the spectral function dominates the long time relaxational dynamics. We find, to the contrary that the Landau damping **thresholds** at $\omega = \pm k$ determine the long- time dynamics and that the early time dynamics is sensitive to several moments of the total spectral density. In ref. [7] we concentrated on the case of scalar electrodynamics (SQED) since this theory has the same HTL structure (to lowest order) as the non-abelian case [11,12]. Most of our results can therefore, be taken over to the non-Abelian case with little or no changes at least to the leading order in HTL.

When finite temperature infrared divergences are present, like for scalar condensates coupled with gauge fields, the amplitude approximation combined with the resummation of the infrared divergences by dynamical renormalization group leads to anomalous logarithmic relaxation and logarithmic phases in time [21].

IV. NON-LINEAR QUANTUM EVOLUTION IN SCALAR FIELD THEORIES

Our goal is to deal with the out of equilibrium evolution for **large** energy densities in field theory. That is, a large number of particles per volume m^{-3} , where m is the typical mass scale in the theory. The most familiar techniques of field theory, based on the S-matrix formulation of transition amplitudes and perturbation theory apply in the opposite limit of low energy density and since they only provide information on *in* $\rightarrow$ *out* matrix elements, are unsuitable for calculations of time dependent expectation values.

Similar tools are also necessary to describe consistently the dynamical processes in the Early Universe [1,3]. In particular it has been recognized that novel phenomena associated with parametric amplification of quantum fluctuations can play an important role in the process of reheating and thermalization [1,3,17]. It must be noticed that the dynamics in cosmological spacetimes

is dramatically different to the dynamics in Minkowski spacetime. Both in fixed FRW and de Sitter [17] backgrounds and in a dynamical geometry [18] the dynamical evolution is qualitatively and quantitatively different to the Minkowski case considered in the present paper.

Our program to study the dynamical aspects of relaxation out of equilibrium both in the linear and non-linear regime has revealed new features of relaxation in the collisionless regime in scalar field theories [3,10]. Recent investigation of scalar field theories in the non-linear regime, including self-consistently the effects of quantum backreaction in an energy conserving and renormalizable framework have pointed out to a wealth of interesting non-perturbative phenomena both in the broken and unbroken symmetry phases [3]. These new phenomena are a consequence of the non-equilibrium evolution of an initial state of large energy density which results in copious particle production leading to non-thermal and non-perturbative distribution of particles. Our studies have focused on the situation in which the *amplitude* of the expectation value of the scalar field is non-perturbatively large, $A \approx \sqrt{\lambda} < \Phi > / m \approx \mathcal{O}(1)$ (m is the mass of the scalar field and λ the self-coupling) and most of the energy of the initial state is stored in the ‘zero mode’, i.e. the (translational invariant) expectation value of the scalar field Φ . Under these circumstances the initial energy density $\varepsilon \approx m^4/\lambda$. During the dynamical evolution the energy initially stored in one (or few) modes of the field is transferred to other modes resulting in copious particle production initially either by parametric amplification of quantum fluctuations in the unbroken symmetry phase, or spinodal instabilities in the broken symmetry phase. Furthermore, we have considered in ref. [14] the evolution of an initial state with a large number of quanta with a distribution around a momentum $|\mathbf{k}_0|$ corresponding to a *thin spherical shell* in momentum space, a ‘tsunami’.

In both cases the mechanism of energy dissipation and particle production results in a number of produced particles per unit volume $\mathcal{N} \propto m^3/\lambda$ which for weak coupling is non-perturbatively large [3,14]. We call ‘linear regime’ to this first stage dominated either by parametric or spinodal instabilities.

We recognized [3] a new *dynamical* time scale t_1 where the linear regime ends. By the time t_1 the effects of the quantum fluctuation on the dynamical evolution become of the same order as the classical contribution given by the evolution of the expectation value of the field. The ‘non-linear’ regime starts by the time t_1 . In the case of broken symmetry, this time scale corresponds to the spinodal scale at which the backreaction of quantum fluctuations shut-off the spinodal instabilities. At this scale non-perturbative physics sets in and the non-linearities of the full quantum theory determine the evolution. This time scale t_1 , which we call the *non-linear time*, is a non-universal feature of the dynamics and depends strongly on the initial state and non-perturbatively on the coupling, as $t_1 \propto \log[\lambda^{-1}]$ for weak coupling [3].

There are two very important parameters that influence the quantum dynamics: the strength of the coupling constant λ and the initial energy density in units of the scalar field mass m . If in the initial state most of the energy is stored in the zero mode, the energy density is determined by the amplitude A of the order parameter. The value of this field amplitude determines the regime of applicability of perturbation theory methods. That is, the amplitude approximation considered in secs. II and III fails to catch important aspects of the physics when A is not small.

We consider now the **non-perturbative** regime in which $A = \sqrt{\lambda} < \Phi > /m \simeq \mathcal{O}(1)$. It is important to point out that for large field amplitude, even for very weakly coupled theories non-linear effects will be important and must be treated non-perturbatively. This is the case under consideration. Having recognized the non-perturbative nature of the problem for large amplitudes we must invoke a non-perturbative, consistent calculational scheme which respects the symmetries (continuous global symmetries and energy-momentum conservation), is renormalizable and lends itself to a numerical treatment.

We are thus led to consider the $O(N)$ vector model with quartic self-interaction [3] and the scalar field in the vector representation of $O(N)$.

The action and Lagrangian density are given by,

$$S = \int d^4x \left\{ \frac{1}{2} [\partial_\mu \Phi(x)]^2 - V(\Phi(x)) \right\},$$

$$V(\Phi) = \frac{\lambda}{8N} \left(\Phi^2 + \frac{2Nm^2}{\lambda} \right)^2 - \frac{Nm^4}{2\lambda}. \quad (30)$$

We write the field Φ as $\Phi = (\sigma, \pi)$ where π represents the $N-1$ ‘pions’, and choose the coupling λ to remain fixed in the large N limit. Due to lack of space we only considered here the unbroken symmetry case $m^2 > 0$. The broken symmetric case ($m^2 < 0$) is considered in ref. [3].

We can decompose the field σ into its expectation value and fluctuations $\chi(\mathbf{x}, t)$ about it:

$$\sigma(\mathbf{x}, t) = \Phi(t) \sqrt{N} + \chi(\mathbf{x}, t), \quad (31)$$

with $\Phi(t)$ being a c-number of order one in the $N \rightarrow \infty$ limit and χ an operator.

We shall not rederive here the field evolution equations for translationally invariant quantum states. In the leading order in the large N approximation the theory becomes Gaussian at the expense of a self-consistent condition [3,15,19], this in turn entails that the Heisenberg field operator $\Psi(\mathbf{x}, t)$ can be written as

$$\Psi(\mathbf{x}, t) = \int \frac{d^3k}{(2\pi)^3} \frac{1}{\sqrt{2}} \left[a_{\mathbf{k}} f_{\mathbf{k}}(t) e^{i\mathbf{k} \cdot \mathbf{x}} + a_{\mathbf{k}}^\dagger f_{\mathbf{k}}^*(t) e^{-i\mathbf{k} \cdot \mathbf{x}} \right],$$

where $a_{\mathbf{k}}, a_{\mathbf{k}}^\dagger$ are the canonical creation and annihilation operators, the mode functions $f_{\mathbf{k}}(t)$ are solutions of the

Heisenberg equations of motion [3,15,19] given below (32-34).

Our choice of initial conditions on the density matrix is that of the vacuum for the instantaneous modes of the Hamiltonian at the initial time [3,19]. Therefore we choose the initial conditions on the mode functions to represent positive energy particle states of the instantaneous Hamiltonian at $t = 0$, which is taken to be the initial time. That is,

$$f_{\mathbf{k}}(0) = \frac{1}{\sqrt{W_{\mathbf{k}}}}; \dot{f}_{\mathbf{k}}(0) = -i\sqrt{W_{\mathbf{k}}} \quad ; \quad W_{\mathbf{k}} = \sqrt{k^2 + M_0^2},$$

where the mass M_0 is defined below. With these boundary conditions, the mode functions $f_{\mathbf{k}}(0)$ correspond to positive frequency modes (particles) of the instantaneous quadratic Hamiltonian for oscillators of mass M_0 .

We point out that the behavior of the system depends mildly on the initial conditions on the mode functions as we have found by varying them within a wide range. In particular, the various types of linear and nonlinear resonances are independent of these initial conditions [3,19].

We introduce the following dimensionless quantities:

$$\tau = |m|t, \quad q = \frac{k}{|m|}, \quad \Omega_q = \frac{W_{\mathbf{k}}}{|m|}, \quad \eta^2(\tau) = \frac{\lambda}{2|m|^2} \Phi^2(t),$$

$$g\Sigma(\tau) = \frac{\lambda}{2|m|^2} [\langle \Psi^2(t) \rangle_R - \langle \Psi^2(0) \rangle_R] \quad , \quad (\Sigma(0) = 0)$$

$$g = \frac{\lambda}{8\pi^2}, \quad \varphi_q(\tau) \equiv \sqrt{|m|} f_{\mathbf{k}}(t).$$

Here $\langle \Psi^2(t) \rangle_R$ stands for the renormalized composite operator [see eq.(35) for an explicit expression].

A. Evolution equations in the large N limit

In terms of the dimensionless variables introduced above the renormalized equations of motion are found to be (see references [3,19]):

$$\ddot{\eta} + \eta + \eta^3 + g\eta(\tau)\Sigma(\tau) = 0 \quad (32)$$

$$\left[\frac{d^2}{d\tau^2} + q^2 + 1 + \eta(\tau)^2 + g\Sigma(\tau) \right] \varphi_q(\tau) = 0, \quad (33)$$

$$\varphi_q(0) = \frac{1}{\sqrt{\Omega_q}}, \quad \dot{\varphi}_q(0) = -i\sqrt{\Omega_q}, \quad \eta(0) = \eta_0, \quad \dot{\eta}(0) = 0 \quad (34)$$

Hence, $\mathcal{M}^2(\tau) \equiv 1 + \eta(\tau)^2 + g\Sigma(\tau)$ plays the rôle of a (time dependent) effective mass squared.

We will choose Ω_q such that at $t = 0$ the quantum fluctuations are in the ground state of the oscillators at the initial time. Since $g\Sigma(0) = 0$, we choose $M_0 \equiv \sqrt{1 + \eta_0^2}$. $g\Sigma(\tau)$ is given by the self-consistent condition [3,19]

$$g\Sigma(\tau) = g \int_0^\infty q^2 dq \left\{ |\varphi_q(\tau)|^2 - \frac{1}{\Omega_q} + \frac{\theta(q-1)}{2q^3} [-\eta_0^2 + \eta^2(\tau) + g\Sigma(\tau)] \right\}. \quad (35)$$

We thus see that the effective mass at time τ contains all q -modes and the zero mode at the same time τ . The evolution equations are then nonlinear but local in time in the infinite N limit.

B. The early time evolution: parametric resonance

In the weak coupling regime the back-reaction term $g\Sigma(\tau)$ is small for small g during an interval say $0 \leq \tau < \tau_1$ [3,19]. This time τ_1 will be called the nonlinear time and it determines the time scale when the backreaction effects and therefore the quantum fluctuations and nonlinearities become important.

For times $\tau < \tau_1$, $g\Sigma(\tau)$ can be neglected and then eq.(32) reduces to the classical equation of motion

$$\ddot{\eta} + \eta + \eta^3 = 0. \quad (36)$$

The solution of this equation with the initial conditions (34) can be written in terms of elliptic functions

$$\eta(\tau) = \eta_0 \operatorname{cn} \left(\tau \sqrt{1 + \eta_0^2}, k \right), \quad k = \frac{\eta_0}{\sqrt{2(1 + \eta_0^2)}},$$

where cn stands for the Jacobi cosine.

Inserting this form for $\eta(\tau)$ in eq.(33) and again neglecting $g\Sigma(\tau)$ yields

$$\left[\frac{d^2}{d\tau^2} + q^2 + 1 + \eta_0^2 \operatorname{cn}^2 \left(\tau \sqrt{1 + \eta_0^2}, k \right) \right] \varphi_q(\tau) = 0. \quad (37)$$

This is the Lamé equation for a particular value of the coefficients that make it solvable in terms of Jacobi functions [3,19].

Since the coefficients of eq.(37) are periodic with period 2ω (notice that $\eta(\tau + 2\omega) = -\eta(\tau)$), the mode functions can be chosen to be quasi-periodic (Floquet type) with quasi-period 2ω ,

$$U_q(\tau + 2\omega) = e^{iF(q)} U_q(\tau), \quad (38)$$

where the Floquet indices $F(q)$ are independent of τ . In the allowed zones, $F(q)$ is real and the functions $U_q(\tau)$ are bounded. In the forbidden zones $F(q)$ has a non-zero imaginary part and the amplitude of the solution either grows or decreases exponentially.

It turns out in the present case that there is only one forbidden band for $q^2 > 0$ running in the interval $0 < q^2 < \eta_0^2/2$.

The modes from the forbidden band $0 < q < \eta_0/\sqrt{2}$ dominate $\Sigma(\tau)$. For $0 < \tau < \tau_1$, $\Sigma(\tau)$ oscillates with an exponentially growing amplitude. This amplitude (envelope) $\Sigma_{env}(\tau)$ can be represented to a very good approximation by the formula [3,19]

$$\Sigma_{env}(\tau) = \frac{1}{N\sqrt{\tau}} e^{B\tau}, \quad (39)$$

where B and N are functions of η_0 given by

$$B(\eta_0) = 8 \sqrt{1 + \eta_0^2} \hat{q} (1 - 4\hat{q}) + O(\hat{q}^3),$$

$$N(\eta_0) = \frac{4}{\sqrt{\pi}} \sqrt{\hat{q}} \frac{(4 + 3\eta_0^2) \sqrt{4 + 5\eta_0^2}}{\eta_0^3 (1 + \eta_0^2)^{3/4}} [1 + O(\hat{q})] \quad (40)$$

and the elliptic nome $\hat{q}$ can be written as a function of η_0 as

$$\hat{q}(\eta_0) = \frac{1}{2} \frac{(1 + \eta_0^2)^{1/4} - (1 + \eta_0^2/2)^{1/4}}{(1 + \eta_0^2)^{1/4} + (1 + \eta_0^2/2)^{1/4}}. \quad (41)$$

with an error smaller than $\sim 10^{-7}$.

Using this estimate for the quantum fluctuations $\Sigma(\tau)$, we estimated the nonlinear time scale τ_1 at which the back-reaction becomes comparable to the classical terms. Such a time is defined by $g\Sigma(\tau_1) \sim (1 + \eta_0^2/2)$, [3]

$$\tau_1 \approx \frac{1}{B(\eta_0)} \log \frac{N(\eta_0) (1 + \eta_0^2/2)}{g \sqrt{B(\eta_0)}}.$$




FIG. 1. The zero mode $\eta(\tau)$ vs. τ for the unbroken symmetry case with $\eta_0 = 4$, $g = 10^{-9}$.

The time interval from $\tau = 0$ to $\tau \sim \tau_1$ is when most of the particle production takes place. Around $\tau \sim \tau_1$ the quantum fluctuation become large enough to begin shutting-off the growth of the modes and particle production slows down dramatically. This dynamical time

scale separates two distinct types of dynamics, for $\tau < \tau_1$ the evolution of the quantum modes $\varphi_q(\tau)$ is essentially linear, the backreaction effects are small and particle production proceed via parametric amplification. For $\tau > \tau_1$ the quantum backreaction effects are as important as the tree level term $\eta(\tau)^2$ and the dynamics is fully non-linear.

The growth of the unstable modes in the forbidden band shows that particles are created copiously ($\sim 1/g$ for $\tau \sim \tau_1$). Initially, ($\tau = 0$) all the energy is in the classical zero mode (expectation value). Part of this energy is rapidly transformed into particles through parametric resonance during the interval $0 < \tau < \tau_1$. At the same time, the field expectation value decreases as is clearly displayed in fig. 1.

The momentum distribution of the particles produced by parametric resonance follows the Floquet index and is peaked at $q \approx \frac{1}{2}\eta_0(1 - \hat{q})$ [3,19]. This is shown in fig. 2.




FIG. 2. Momentum distribution of the produced particles $n(q) \equiv q^2 N_q^{ad}(\tau)$ at the nonlinear time $\tau = \tau_1$ for $\eta_0 = 4$, $g = 10^{-9}$.

For late times, typically larger than 3 or 4 times τ_1 , the effective mass tends asymptotically to a constant value $\mathcal{M}_\infty^2 = 1 + \frac{\eta_0^2}{2}$. This fact implies that the modes becomes effectively **free**.

At the same time, the field expectation value relaxes to zero asymptotically with a non-universal power law. The initial energy density which is non-perturbatively large goes completely into production of massive particles. The asymptotic particle distribution is localized within a band determined by the initial conditions with

non-perturbatively large amplitude $\sim 1/\sqrt{\lambda}$ which could be described as a ‘semiclassical condensate’ in the unbroken phase [see fig. 3].




FIG. 3. Momentum distribution of the produced particles at $\tau = 3000$ for $\eta_0 = 1$, $g = 10^{-3}$. Notice the main peak at $q = 0.067$, in good agreement with the estimate K_1/τ given in ref.[2].

The novel result that emerged combining numerical methods and dynamical renormalization group analysis is the presence of *non-linear resonances* that lead to asymptotic relaxation described by *non-universal power laws* [3]. These power laws are determined by *dynamical anomalous exponents* which depend non-perturbatively on the coupling.

V. ACKNOWLEDGEMENTS

D. B. thanks the N.S.F. for partial support through grant PHY-9605186. R. H. and S. P. K. were supported by DOE grant DE-FG02-91-ER40682. The authors acknowledge support from NATO. H. J. de V. thanks the organizers of TFT98 where this work has been reported.

- [1] See for example, D. Boyanovsky, D. Cormier, H. J. de Vega, R. Holman and S. Prem Kumar, Erice Chalonge School, Current Topics in Astrofundamental Physics, N. Sánchez and A. Zichichi Editors, p. 119-194, Kluwer, 1998 and references therein.
- [2] J. Schwinger, J. Math. Phys. 2, 407 (1961); K. T. Mahanthappa, Phys. Rev. 126, 329 (1962); P. M. Bakshi and K. T. Mahanthappa, J. Math. Phys. 41, 12 (1963); L. V. Keldysh, JETP 20, 1018 (1965); K. Chou, Z. Su, B. Hao And L. Yu, Phys. Rep. 118, 1 (1985); A. Niemi and G. Semenoff, Ann. of Phys. (NY) 152, 105 (1984); N. P. Landsmann and C. G. van Weert, Phys. Rep. 145, 141 (1987); E. Calzetta and B. L. Hu, Phys. Rev. D41, 495 (1990); *ibid* D37, 2838 (1990); J. P. Paz, Phys. Rev. D41, 1054 (1990); *ibid* D42, 529(1990).
- [3] D. Boyanovsky, H. J. de Vega and R. Holman, Proceedings of the Second Paris Cosmology Colloquium, Observatoire de Paris, June 1994, pp. 127-215, H. J. de Vega and N. Sanchez, Editors (World Scientific, 1995); Advances in Astrofundamental Physics, Erice Chalonge Course, N. Sanchez and A. Zichichi Editors, (World Scientific, 1995); D. Boyanovsky, H. J. de Vega, R. Holman, D.-S. Lee and A. Singh, Phys. Rev. **D51**, 4419 (1995); D. Boyanovsky, H. J. de Vega, R. Holman and J. Salgado, Phys. Rev. **D54**, 7570 (1996). D. Boyanovsky, C. Destri, H. J. de Vega, R. Holman et J. F. J. Salgado, Phys. Rev. **D57**, 7388 (1998).
- [4] J. W. Harris and B. Muller, Annu. Rev. Nucl. Part. Sci. 46, 71 (1996). B. Muller in *Particle Production in Highly Excited Matter*, Eds. H.H. Gutbrod and J. Rafelski, NATO ASI series B, vol. 303 (1993). B. Muller, *The Physics of the Quark Gluon Plasma* Lecture Notes in Physics, Vol. 225 (Springer-Verlag, Berlin, Heidelberg, 1985); B. Muller, *The Physics of the Quark Gluon Plasma* Lecture Notes in Physics, Vol. 225 (Springer-Verlag, Berlin, Heidelberg, 1985). K. Geiger, Phys. Rep. 258, 237 (1995); Phys. Rev. D46, 4965 (1992); Phys. Rev. D47, 133 (1993); *Quark Gluon Plasma 2*, Ed. by R. C. Hwa, (World Scientific, Singapore, 1995). X. N. Wang, Phys. Rep. 280, 287 (1997). M. H. Thoma, in *Quark Gluon Plasma 2*, ed. by R. C. Hwa, (World Scientific, Singapore, 1995).
- [5] J.-P. Blaizot, J. -Y. Ollitrault and E. Iancu, in *Quark Gluon Plasma 2*, Ed. by R. C. Hwa, (World Scientific, Singapore, 1995).
- [6] H. Elze and U. Heinz, Phys. Rep. 183, 81 (1989).
- [7] D. Boyanovsky, H. J. de Vega, S. P. Kumar and R. Pisarski, 'Real-time relaxation and kinetics in hot scalar QED: Landau damping', hep-ph/9802370, to appear in Phys. Rev. D.
- [8] R. D. Pisarski, Physica A158, 146 (1989); Phys. Rev. Lett. 63, 1129 (1989). E. Braaten and R. D. Pisarski, Nucl. Phys. B337, 569 (1990); E. Braaten and R. D. Pisarski, Nucl. Phys. B339, 310 (1990). R. D. Pisarski, Nucl. Phys. B309, 476 (1988). J. Frenkel and J. C. Taylor, Nucl. Phys. B334, 199 (1990); J. C. Taylor and S. M. H. Wong, Nucl. Phys. B346, 115 (1990); R. Efraty and V. P. Nair, Phys. Rev. Lett. 68, 2891 (1992), R. Jackiw and V. P. Nair, Phys. Rev. D48, 4991 (1993). J. P. Blaizot and E. Iancu, Nucl. Phys. B421, 565, (1994); Nucl. Phys. B434, 662, (1995); Phys. Rev. Lett. 70, 3376 (1993).
- [9] J.-P. Blaizot, E. Iancu, Phys.Rev. D55 973 (1997) and D52, 254 (1995) ; Phys.Rev.Lett. 76 (1996); Nucl.Phys. B459, 559 (1996).
- [10] D. Boyanovsky, M. D'Attanasio, H.J. de Vega and R. Holman, Phys. Rev. **D54**, 1748 (1996).
- [11] M. Le Bellac, Thermal Field Theory, (Cambridge University Press, 1996).
- [12] U. Kraemmer, A.K.Rebhan, and H. Schulz, Annals Phys. 238, 286 (1995).
- [13] D. Boyanovsky, H. J. de Vega and R. Holman, Phys. Rev. **D 51**, 734 (1995). D. Boyanovsky, H. J. de Vega, R. Holman and S. Prem Kumar, Phys. Rev. **D56**, 3929 and 5233 (1997).
- [14] D. Boyanovsky, H. J. de Vega, R. Holman, S. Prem Kumar and R. D. Pisarski, Phys. Rev. **D57**, 3653 (1998).
- [15] Y. Kluger, J. M. Eisenberg, B. Svetitsky, F. Cooper and E. Mottola, Phys. Rev. Lett. 67, 2427 (1991); Phys. Rev. D 45, 4659, (1992); Phys. Rev. D48, 190 (1993); F. Cooper, S. Habib, Y. Kluger, E. Mottola, J. P. Paz, P. R. Anderson, Phys. Rev. D50, 2848 (1994). F. Cooper, S.-Y. Pi and P. N. Stancioff, Phys. Rev. D34, 3831 (1986). F. Cooper, Y. Kluger, E. Mottola, J. P. Paz, Phys. Rev. D51, 2377 (1995). F. Cooper and E. Mottola, Phys. Rev. D36, 3114 (1987).
- [16] D. Boyanovsky, H. J. de Vega and R. Holman, Vth. Erice Chalonge School, Current Topics in Astrofundamental Physics, N. Sánchez and A. Zichichi Editors, World Scientific, 1996, p. 183-270. D. Boyanovsky, M. D'Attanasio, H. J. de Vega, R. Holman and D.-S. Lee, Phys. Rev. **D52**, 6805 (1995).
- [17] D. Boyanovsky, D. Cormier, H.J. de Vega, R. Holman, A. Singh, M. Srednicki, Phys. Rev. **D56**, 1939 (1997). D. Boyanovsky, D. Cormier, H.J. de Vega and R. Holman, Phys. Rev **D55**, 3373 (1997).
- [18] D. Boyanovsky, D. Cormier, H. J. de Vega, R. Holman and S. P. Kumar, Phys. Rev. **D57**, 2166, (1998).
- [19] D. Boyanovsky, H. J. de Vega and R. Holman, p. 183-270 in the Proceedings of the Vth. Erice School 'D. Chalonge', 'Current Topics in Astrofundamental Physics', N. Sánchez and A. Zichichi, editors, World Scientific.
- [20] A. Erdélyi et al., Higher Transcendental Functions, vol. III, p. 217, Mc Graw Hill, New York, 1955.
- [21] D. Boyanovsky, H. J. de Vega, R. Holman and M. Simionato, 'Dynamical Renormalization Group Resummation of Finite Temperature Infrared Divergences', hep-ph/9809346.